arXiv:1210.4267v2 [cond-mat.dis-nn] 9 Nov 2012

Complete absence of localization in a family of disordered lattices

Biplab Pal,[1, *] Santanu K. Maiti,[2, †] and Arunava Chakrabarti[1, ‡]
[1] *Department of Physics, University of Kalyani, Kalyani, West Bengal-741 235, India*
[2] *Physics and Applied Mathematics Unit, Indian Statistical Institute,*
203 Barrackpore Trunk Road, Kolkata-700 108, India

We present analytical results supported by numerical analysis to show that even in a discrete tight-binding model, a random arrangement of a simple four-atom diamond cell in a host lattice of identical atoms can render the entire energy spectrum absolutely continuous when special correlations in the *numerical values* of a subset of the system parameters are enforced. The observation is in total contrast to the canonical case of Anderson localization, which is insensitive to the numerical values of the parameters describing the system. The result has been verified for a wide variety of geometries, and rings true in every case.

PACS numbers: 71.30.+h, 72.15.Rn, 03.75.-b

Localization of electronic states in a disordered lattice, first proposed by Anderson [1] is a problem of everlasting interest in condensed matter physics, and continues to generate intriguing features in quantum transport properties of randomly disordered systems. Over the years, with the improvement of fabrication and lithographic techniques the realm of Anderson localization [2, 3] has extended beyond the electronic systems, and has encompassed tailor made artificial crystal structures, viz., the photonic [4, 5], phononic [6, 7], plasmonic [8, 9] or polaritonic [10, 11] lattices. Very recently, ultracold gases even allowed for the direct observation of localization of matter waves [12–14].

The pivotal result in this field is that, the electronic wave functions are known to be localized for dimensions $d \leq 2$ (the band center in the off diagonal disorder case is an exception), and even for $d > 2$ for strong disorder, with an exponential decay in the envelope of the wave functions [2, 3]. The result has been substantiated by meticulous analyses of various calculations related to the localization length [15, 16], density of states [17], or multi-fractality of the single particles states [18, 19]. Extensive work has also been undertaken to study the intricacies of the single parameter scaling hypothesis - its validity [20], variance [21], or even violation [22, 23] in low dimensional systems within a tight-binding approximation, that has been subsequently consolidated by experimental measurements of conductance distribution in quasi-one dimensional gold wires [24].

In the last decade we have come across examples, within a tight-binding description, where localization-delocalization transitions have been observed in disordered systems. The transitions are attributed to certain special kinds of correlation in the potential profiles [25–28] and, have unraveled the presence of discrete energy levels corresponding to *extended eigenfunctions* [25]. This led to the possibility of a spectral continuum and metal-insulator transition [26, 27] in one, or quasi-one dimensional discrete systems. The idea of engineering extended states in two dimensional disordered systems [28] has also been put forward recently. In such cases the general exponentially localized character of the eigenfunctions prevail, and there is a mixed spectrum of localized and extended states (under some special correlations as discussed above).

In the present letter, we put forward an example where the fundamental result of Anderson localization seems to get violated. We consider a diamond network of four atomic like sites placed at random in a host lattice of isolated atomic sites. The system is described by a tight-binding Hamiltonian. The on-site potential at every lattice point is taken to be a constant, while the nearest-neighbor hopping integral along the arms of the diamond (λ) differ from that along the back bone (t), where it is assumed to be constant (see Fig. 1). There is no positional correlation, short range or long range in the conventional sense [25, 26]. We show that, for a particular relationship between the nearest-neighbor hopping integrals, the infinite *diamond-dot* (DD) chain will yield an absolutely continuous spectrum, and that, the two terminal transmission coefficient will be unity, irrespective of the energy of the electron. This result is true not only for such a DD chain, but also for other geometrical arrangements of the



FIG. 1: (Color online). (a) A typical realization of an array of isolated atomic sites (marked as black circles) and the diamond shaped quadruplets (in isolation, and in clusters, and marked by light blue circles. (b) The renormalized 1-d chain, where the isolated dots remain un-renormalized, while the diamond clusters are renormalized into β-γ doublets, or a sequence of β-δ^n-γ. The hopping integral along the major axis (backbone) is t, while it is λ along the edges of a diamond. All on-site potentials are equal.

side coupled atomic clusters with respect to the linear chain backbone. We explain in details the DD results in this letter with a brief mention of the other cases that we have come across. The results, as they appear, are in complete contradiction to the canonical case of Anderson localization with any kind of disorder, where localization does not depend on the numerical values assigned to the system parameters (and is thus *universal*). To the best of our knowledge, such examples have never been discussed in the literature so far.

Spinless, non-interacting electrons on the DD chain are described by the Hamiltonian,

$$H_{DD} = \epsilon \sum_i d_i^\dagger d_i + \sum_{\langle ij \rangle} t_{ij} \left[d_i^\dagger d_j + h.c. \right] \qquad (1)$$

where, ϵ is the constant on-site potential, the nearest-neighbor hopping integral $t_{ij} = t$ along the backbone, and $t_{ij} = \lambda$ along an edge of the diamond. The Schrödinger equation, written equivalently in the form of the difference equation, $(E-\epsilon)\psi_i = \sum_j t_{ij}\psi_j$ allows us to decimate out the vertices of the diamond networks to map the original chain on to an effective one dimensional chain [Fig. 1(b)] of four sites α, β, γ and δ with on-site potentials $\epsilon_\alpha = \epsilon$, $\epsilon_\beta = \epsilon_\gamma = \epsilon + 2\lambda^2/(E-\epsilon)$, and $\epsilon_\delta = \epsilon + 4\lambda^2/(E-\epsilon)$. There is a binary distribution of hopping integrals along the chain now. These are the original t, and the effective coupling $\tau = 2\lambda^2/(E-\epsilon)$ arising out of the renormalization of the diamond network.



FIG. 2: (Color online). (a) A perfectly periodic array of identical atomic sites (dots). (b) The renormalized lattice obtained by decimating the encircled dots in (a). This generates a chain where the β, γ and δ sites and the nearest-neighbor hopping integrals have precisely those values as obtained in the disordered DD array by setting $\lambda = t/\sqrt{2}$.

Using the difference equation the amplitudes of the wave function at the neighboring sites along the effective one dimensional chain can be related using the 2×2 transfer matrices,

$$\begin{pmatrix} \psi_{n+1} \\ \psi_n \end{pmatrix} = \begin{pmatrix} \dfrac{E-\epsilon_n}{t_{n,n+1}} & -\dfrac{t_{n,n-1}}{t_{n,n+1}} \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \psi_n \\ \psi_{n-1} \end{pmatrix} \qquad (2)$$

A look at the Fig. 1(b) will make it obvious that there are four kinds of transfer matrices, viz, M_α, M_β, M_γ and M_δ, which will differ in their matrix elements, depending on the respective on-site potentials and the nearest-neighbor hopping integrals. From the arrangement of the diamonds and the isolated sites in the original DD chain it can be appreciated that the the wave function at a far end of the chain can be determined if one evaluates the product of the unimodular matrices M_α, $M_{\gamma\beta} = M_\gamma . M_\beta$ and $M_{\gamma\delta\beta} = M_\gamma . M_\delta . M_\beta$, or $M_{\gamma\delta^n\beta} = M_\gamma . M_\delta^n . M_\beta$ sequenced in the desired random fashion.

The central result of this communication is that, the commutators $[M_\alpha, M_{\beta\gamma}] = 0$, and $[M_\alpha, M_{\gamma\delta\beta}] = 0$ irrespective of the energy E of the electron whenever we choose $\lambda = t/\sqrt{2}$. In an infinite array of diamonds and dots, without any restriction on randomness, there can be isolated diamonds (equivalent to a β-γ pair), or a sequence of $n+1$ diamonds (equivalent to a sequence of clusters β-δ^n-γ). The total transfer matrix for such a cluster can be written as, $M_{\gamma\delta^n\beta} = U_{n-1}(x) M_{\gamma\delta\beta} - U_{n-2}(x) M_{\gamma\beta}$, where, $U_n(x)$ is the n-th order Chebyshev polynomial of the second kind and $x = Tr(M_\delta)/2$. It immediately becomes obvious that $[M_\alpha, M_{\gamma\delta^n\beta}] = 0$ for any value of n. This implies that, with $\lambda = t/\sqrt{2}$ the amplitude of the wave function or its phase at any lattice point on the *renormalized* chain [Fig. 1(b)] in the actual randomly disordered array of diamond and dots will be indistinguishable from that *in a perfectly periodic arrangement of these clusters*. This happens independent of the energy E of the electron, that is, throughout the energy spectrum, and is true for any kind of disordered arrangement that one can build using a dot, isolated diamonds or an array of any n number of diamonds. The wave functions as result, will have to be of a *perfectly*



FIG. 3: (Color online). Transmission probability T (red color) and Average Density of States (ADOS) (green color) as a function of energy E for a completely disordered array of diamonds and dots in equal concentration. The total number of sites is 600, and results averaged over 50 disorder configurations have been presented. In the non-resonant case (a) we have chosen $\lambda = 0.3$, while $\lambda = 1/\sqrt{2}$ in (b) where we have complete transparency. Other parameters are, $t = 1 = t_0$, $\epsilon_i = 0$ at every site, $\tau_L = \tau_S = 1$, and $\epsilon_0 = 0$ in the leads.

extended, Bloch-like character.

The energy spectrum in the above case will have to be *absolutely continuous*, and the local density of states (LDOS) at any nodal point α, β, γ or δ on the renormalized lattice will resemble that for a perfectly ordered lattice of identical sites with on-site potential ϵ and nearest neighbor hopping integral t. The band extends from $[\epsilon - 2t, \epsilon + 2t]$. We have extensively verified this, though present only the average density of states to save space. Nevertheless, the fact that it should be the reality, can be tested by observing that under the condition $\lambda = t/\sqrt{2}$, the on-site potentials on the renormalized chain in Fig. 1(b) are, $\epsilon_\alpha = \epsilon$, $\epsilon_\beta = \epsilon_\gamma = \epsilon + t^2/(E-\epsilon)$, and $\epsilon_\delta = \epsilon + 2t^2/(E-\epsilon)$. The nearest-neighbor hopping matrix elements turn out to be t (unchanged value) and $\tau = t^2/(E-\epsilon)$. Most interestingly, identical values can be obtained by beginning with a perfectly periodic chain with constant on-site potential ϵ and nearest-neighbor hopping t, and by arbitrarily decimating sites so as to reproduce the same disordered pattern as the original chain. The process is illustrated in Fig. 2. Now, the lattice in Fig. 2 is a perfectly periodic one, and no matter how we decimate sites, the LDOS at any site will be that of a periodic chain with van Hove singularities marking the band edges. This is what makes the LDOS at any site on the backbone of our original system in Fig. 1(b) indistinguishable from that of an ordered chain of atoms.



FIG. 4: (Color online). Transmission probability T (red color) and ADOS (green color) as a function of energy E for the diamond-dot geometry with a total of 600 sites and for arbitrary concentration. Results averaged over 50 different configurations have been presented. Again we have selected $t = 1$ and $\lambda = 0.3$ for the non-resonant case (a), and $\lambda = 1/\sqrt{2}$ in the resonant case (b). Other parameters are the same as in Fig. 3.

To obtain the energy spectrum and the transmission probability we use a Green's function formalism [29]. Keeping in mind a possible experimental realization of the system, we clamp a finite sized system of N-sites between two ideal semi-infinite electrodes (the left and the right electrodes) making an electrode-system-electrode (ESE) bridge. The electrodes are described by the Hamiltonian,

$$H_0 = \sum_i \epsilon_0 c_i^\dagger c_i + \sum_{<ij>} t_0 \left(c_i^\dagger c_j + c_j^\dagger c_i \right) \tag{3}$$

where different parameters correspond to their usual meaning. The couplings between the left (L) and the right (R) electrodes and the system (S) are given by, $H_{LS} = \tau_L c_1^\dagger d_1 + h.c.$, and $H_{RS} = \tau_R d_N^\dagger c_1 + h.c.$, so that, the full Hamiltonian of the ESE bridge is given by, $H = H_0 + H_{LS} + H_{RS} + H_{DD}$. The average density of states (ADOS) of the ESE bridge is given by,

$$\rho_{av} = -\frac{1}{N\pi} \text{Im}\left[\text{Tr}(G)\right] \tag{4}$$

where, $G = (E - H + i\eta)^{-1}$, with $\eta \to 0$.

In terms of the Green's function of the conductor and its coupling to the side-attached electrodes, the transmission probability can be written in the form [29],



FIG. 5: (Color online). Transmission probability T (red color) and ADOS (green color) as a function of energy E for the diamond-dot chain away from the resonant case. The results averaged over 50 disorder configurations have been presented. 100 diamonds with 100 dots have been taken with $\lambda = 0.98499$, a forty percent deviation from the resonance value of $1/\sqrt{2}$. Other parameters are the same as in the earlier figures.

$$T = \text{Tr}\left[\Gamma_L \, G_c^r \, \Gamma_R \, G_c^a\right] \tag{5}$$

where, Γ_L and Γ_R describe the coupling of the conductor to the left and right electrodes, respectively. Here, G_c^r and G_c^a are the retarded and advanced Green's functions, respectively, of the conductor including the effects of the electrodes.

Evaluation of this Green's function, both for getting the ADOS and the transmittance needs the inversion of an infinite matrix, which is an impossible task, since the full system consists of the finite size sample and two semi-infinite 1D leads. However, the full system can be partitioned into sub-matrices corresponding to the individual sub-systems and the Green's function for the disordered sample can be effectively written as,

$$G_c = (E - H_c - \Sigma_L - \Sigma_R)^{-1} \tag{6}$$

where, Σ_L and Σ_R are the self-energies due to coupling of the conductor to the left and right electrodes, respectively [29]. All information of the coupling are included into these two self-energies.

In Fig. 3 and Fig. 4 we show the ADOS and the transmission coefficient of a DD sequence of a total of 600 atomic sites. While, in the former, the diamonds and the dots appear in equal concentration, the distribution is completely unrestricted in the latter. Our previous analytical argument for the infinite system is perfectly corroborated in these diagrams. In both the figures the complete localization of all the eigenstates, corresponding to zero transmission, are observed [Fig. 3(a) and Fig. 4(a)] as long as λ is set quite arbitrarily. The scenario changes dramatically as soon as the value of λ is set equal to the calculated value of $1/\sqrt{2}$ (with $t = 1$). We now see an absolutely continuous energy spectrum between $E = \pm 2t$, the central peak at $E = 0$ being the contributions from the top and bottom sites of the diamond plaquettes, and the transmission coefficient becomes equal to unity *irrespective of the energy* of the electron.



FIG. 6: (Color online). A disordered arrangement of triangular plaquettes threaded by a magnetic flux, and dots. The lattice is completely transparent to an incoming electron ($-2t < E < 2t$) if one chooses $\lambda = t/\sqrt{2}$ and $\Phi = \Phi_0/4$.

Once again, keeping in mind a possible experimental realization of such a finite size sample, we have carefully studied the effect of a possible deviation in the value of λ from its ideal 'resonance value' of $1/\sqrt{2}$. The continuity of the spectrum stands out to be a robust result even when the deviation is substantial. In Fig. 5 we show the ADOS and the transmission coefficient for a 40 % deviation in the value of λ. The transmission spectrum is seen to be split into three continuous sub-bands of high transmission. Interestingly, the transmission spectrum is seen to be zero just outside a continuous high transmission zone where the ADOS is still non zero. The system, with such parameter setting thus opens up the possibility of exhibiting even a metal-insulator transition, but this needs much more careful investigation.

Before we end, it may be pointed out that a family of similar disordered quasi-one dimensional array of scatterers can be made to exhibit an absolutely continuous spectrum even for any disordered arrangement of the basic building blocks. Figure 6 refers to one such case where a random arrangement of triangular plaquettes threaded by a magnetic flux Φ and dots are presented. For an identical value of the on-site potential at every site, this system becomes completely transparent to an incoming electron of any energy $-2t < E < 2t$ provided we set $\lambda = t/\sqrt{2}$ and $\Phi = \Phi_0/4$. Even simpler geometries with a minimal side coupling of one quantum dot is capable of sustaining an absolutely continuous spectrum in a lattice with off diagonal disorder [30]. Details of these cases will be presented elsewhere.

Biplab Pal would like to thank DST, India for providing financial assistance through an INSPIRE Fellowship (IF110078). Santanu K. Maiti is thankful to Abraham Nitzan for stimulating discussions, and gratefully acknowledges the hospitality at the School of Chemistry, Tel Aviv University where part of the computation was done.

---

[*] Electronic address: biplabpal@klyuniv.ac.in
[†] Electronic address: santanu.maiti@isical.ac.in
[‡] Electronic address: arunava@klyuniv.ac.in

[1] P. W. Anderson, Phys. Rev. **109**, 1492 (1958).
[2] B. Kramer and A. MacKinnon, Rep. Prog. Phys. **56**, 1469 (1993).
[3] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
[4] E. Yablonovitch, Phys. Rev. Lett. **58**, 2059 (1987).
[5] S. John, Phys. Rev. Lett. **58**, 2486 (1987).
[6] F. R. Montero de Espinosa, E. Jiménez, and M. Torres, Phys. Rev. Lett. **80**, 1208 (1998).
[7] J. O. Vasseur, P. A. Deymier, G. Frantziskonis, G. Hong, B. Djafari-Rouhani, and L. Dobrzynski, J. Phys.: Condens. Matter **10**, 6051 (1998).
[8] A. Tao, P. Sinsermsuksaul, and P. Yang, Nature Nanotechnology **2**, 435 (2007).
[9] A. Christ, Y. Ekinci, H. H. Solak, N. A. Gippius, S. G. Tikhodeev, and O. J. F. Martin, Phys. Rev. B **76**, 201405(R) (2007).
[10] I. O. Barinov, A. P. Alodzhants, and S. M. Arakelian, Quantum Electronics **39**, 685 (2009).
[11] M. Grochol and C. Piermarocchi, Phys. Rev. B **78**, 035323 (2008).
[12] B. Damski, J. Zakrzewski, L. Santos, P. Zoller, and M. Lewenstein, Phys. Rev. Lett. **91**, 080403 (2003).
[13] J. Billy, V. Josse, Z. Zuo, A. Bernard, B. Hambrecht, P. Lugan, D. Clément, L. Sanchez-Palencia, P. Bouyer, and A. Aspect, Nature (London) **453**, 891 (2008).
[14] G. Roati, C. D'Errico, L. Fallani, M. Fattori, C. Fort, M. Zaccanti, G. Modugno, M. Modugno, and M. Inguscio, Nature (London) **453**, 895 (2008).
[15] R. A. Römer and H. Schulz-Baldes, Europhys. Lett. **68**, 247 (2004).
[16] A. Eilmes, R. A. Römer, and M. Schreiber, Physica B **296**, 46 (2001).
[17] A. Rodríguez, J. Phys. A: Math. Gen. **39**, 14303 (2006).
[18] A. Rodriguez, L. J. Vasquez, and R. A. Römer, Phys. Rev. B **78**, 195107 (2008).
[19] A. Rodriguez, L. J. Vasquez, K. Slevin, and R. A. Römer, Phys. Rev. B **84**, 134209 (2011).
[20] S. D. Pinski, W. Schirmacher, and R. A. Römer, Euro-

phys. Lett. **97**, 16007 (2012).

[21] Lev I. Deych, A. A. Lisyansky, and B. L. Altshuler, Phys. Rev. Lett. **84**, 2678 (2000).

[22] J. W. Kantelhardt and A. Bunde, Phys. Rev. B **66**, 035118 (2002).

[23] M. Titov and H. Schomerus, Phys. Rev. Lett. **95**, 126602 (2005).

[24] P. Mohanty and R. A. Webb, Phys. Rev. Lett. **88**, 146601 (2002).

[25] D. H. Dunlap, H-L. Wu, and P. W. Phillips, Phys. Rev. Lett. **65**, 88 (1990).

[26] F. A. B. F. de Moura and M. L. Lyra, Phys. Rev. Lett. **81**, 3735 (1998).

[27] S. Sil, S. K. Maiti, and A. Chakrabarti, Phys. Rev. B **78**, 113103 (2008).

[28] A. Rodriguez, A. Chakrabarti, and R. A. Römer, Phys. Rev. B **86**, 085119 (2012).

[29] S. Datta, *Electronic transport in mesoscopic systems*, Cambridge University Press, Cambridge (1995).

[30] A. Chakrabarti and S. Chattopadhyay, e-print arXiv:1112.0871.